

Inelastic J/ψ electroproduction in e-p collisions

P. Balbi, A. Giovannini
Theoretical Physics Department, University of Torino

February 1, 2008

Report-no: DFTT 39/02

Following HERA experiments our interest is focused, in the present paper, on charmonium production in lepton-proton collisions.

Inelastic J/ψ electroproduction is quite difficult to be measured due to the low rates of events; almost all experimental studies are indeed performed in photoproduction regime.

J/ψ resonance is considered as the product of the hadronization of a $c\bar{c}$ pair generated via boson-gluon fusion.

The leading order contribution to the differential cross section, with respect to virtual photon-proton center of mass energy W and transferred four-momentum squared Q^2 , is written as the sum of two terms related to a colour singlet and a colour octet $c\bar{c}$ pair production respectively.

This expression contains two-gluons contributions and depends on three parameters; two of them are related to the hadronization fractions in charmonium states for the $c\bar{c}$ pair (colour singlet and colour octet), while the third parameter takes into account possible “hot-spot” effects.

The values of these parameters are estimated by mean of a fit to the experimental data collected by H1 collaboration at HERA.

By using different gluon densities, different sets of parameters are obtained, always with good fits.

Hidden charm production in e-p collisions at HERA is an intriguing subject both from the theoretical and from experimental point of view.

Available experimental data concern mainly charmonium photoproduction which corresponds to $Q^2 \rightarrow 0$ limit ($q^2 = -Q^2$ is the transferred four-momentum squared) [1] [2] [3] [4].

In view of the low rates of events (cross sections decreasing with respect to Q^2) only one sample of data (taken by H1 detector) on J/ψ electroproduction is available [5].

It should be pointed out that $e p \rightarrow e J/\psi X$ photoproduction cross sections are characterized by one perturbative mass scale, namely the charm quark mass m_c .

These cross sections are found to be dominated by diffractive processes whose hallmark is the low value for X mass M_X .

Coming to electroproduction events there are two perturbative mass scales which come into play, m_c and Q .

Either diffractive photoproduction or electroproduction experimental rates have been recently understood in the framework of perturbative QCD (pQCD) [6] [7] [8].

The main aim of this paper is the analysis of inelastic J/ψ electroproduction; the true inelastic events are characterized by a M_X value much higher than the diffractive ones.

The inelastic contributions to the inclusive cross sections can be therefore isolated by mean of a cutoff M_{min} on the values for M_X (H1 coll. suggests $M_{min} = 10 \frac{GeV}{c^2}$ [5]).

H1 experimental points cover the kinematical range

$$2 < Q^2 < 80 \text{ GeV}^2$$

so that electroweak corrections (Z_0 exchange) can be neglected and one can work in single-photon-exchange approximation.

HERA results on open charm production in e^+p DIS suggest to describe also inelastic charmonium electroproduction in pQCD by $c\bar{c}$ pairs formation via boson-gluon fusion (BGF) mechanism given by the following partonic subprocess :

$$\gamma^* g \rightarrow c \bar{c} X$$

That is to say any intrinsic charm contribution to the proton wave-

function is neglected.




Figure 1: BGF mechanism

The Born contribution is given by the following partonic diagrams:




Figure 2: Born terms for $e g \rightarrow e c \bar{c}$

Of course these diagrams contribute to the cross sections for production of a $c\bar{c}$ pair in a colour octet state (this pair is coupled to a gluon-photon system)

$$e p \rightarrow e |c\bar{c}\rangle_8 X$$

The emission of an additional gluon allows the $c\bar{c}$ pair to emerge in a colour singlet state; so the lowest order contribution to the

$$e p \rightarrow e |c\bar{c}\rangle_1 X$$

process comes from the following diagrams




Figure 3: Lowest order contributions to $e g \rightarrow e g c \bar{c}$

Diagrams in Figure 3 contribute both to $e p \rightarrow e |c\bar{c}\rangle_1 X$ and to $e p \rightarrow e |c\bar{c}\rangle_8 X$.

It should be pointed out that the two other diagrams shown in figure 4 contribute to $e p \rightarrow e |c\bar{c}\rangle_8 X$ process at the same perturbative order.




Figure 4: Lowest order contributions to $e g \rightarrow e g |c\bar{c}\rangle_8$

If we examine the overall lowest order contribution to the squared matrix element for $e g \rightarrow e g c \bar{c}$ transition, we easily find that it enjoys the following colour decomposition:

$$\overline{\sum} |M_{LO}(e g \rightarrow e g c \bar{c})|^2 = S_{QED} T(F) C_2(F) + S_{NA} C_2(A) \quad (1)$$

where the symbol $\overline{\sum}$ means that we are averaging over colour and polarization indexes for incoming particles and summing over outgoing states.

S_{QED} and S_{NA} are the “space-time” coefficients; S_{QED} takes contribution only from the diagrams in figure 3, while S_{NA} includes also diagrams in figure 4.

The colour factors are given, as usual, by

$$C_2(F) = \frac{N^2 - 1}{2N} = \frac{4}{3}, \quad T(F) = \frac{1}{2}, \quad C_2(A) = N = 3$$

The lowest order terms of the gluon initiated cross sections for a colour singlet $c\bar{c}$ pair inclusive production $\sigma_g(e p \rightarrow e |c\bar{c}\rangle_1 X)$ turn out to be collinear finite since the charm quark is massive.

Moreover the soft divergences found in the perturbative expansions for $\sigma_g(e p \rightarrow e |c\bar{c}\rangle_8 X)$ and $\sigma_g(e p \rightarrow e |c\bar{c}\rangle_1 X)$ compensate each other (order by order in perturbation theory); then the cross sections

$\sigma_g(e p \rightarrow e c \bar{c} X)$ summed over colour states for the $c\bar{c}$ pair remain finite. They have been explicitly computed in the literature, up to next to leading order [9] [10] [11].

If we consider the colour structure of the lowest order squared matrix element for the transition

$$e g \rightarrow e |c\bar{c}\rangle_1 g$$

we find it's given by

$$\sum |M_{LO}(e g \rightarrow e |c\bar{c}\rangle_1 g)|^2 = S_{QED} C_1, \quad C_1 = \frac{1}{8} T(F) C_2(F)$$

where C_1 is the corresponding colour factor.

It's important to notice that H1 events run over the ranges

$$40 < W < 180 \text{ GeV}, 0.2 < z \lesssim 1$$

where W is the center of mass energy of the virtual photon-proton system and z is the ratio of J/ψ and exchanged photon energies in the proton rest frame (elasticity of the process).

It can be easily shown that the Bjorken scaling variable x , in terms of W and Q^2 , is given by $x = \frac{Q^2}{W^2 + Q^2}$.

Accordingly the experimental points lie in the low x region

$$6 \cdot 10^{-5} < x < 5 \cdot 10^{-2}$$

It's well known that the density of gluons $\rho_g(\xi, \mu)$, with respect to the fraction of longitudinal momentum carried (ξ) and the factorization scale (μ), shows a rapid enhancement as ξ decreases.

For x values small enough gluon recombination effects can no longer be neglected; in other words the BGF dynamics could involve 2 gluons (boson-gluon-gluon fusion).

It should be stressed that these effects contribute to the cross sections for inclusive colour singlet $|c\bar{c}\rangle_1$ pair production already at the next-to-leading order; so we have explicitly calculated (under appropriate hypotheses) the lowest order contribution of the

$$e g g \rightarrow e |c\bar{c}\rangle_1$$

partonic subprocess to the double-differential cross section with respect to W and Q^2 variables

$$\frac{d^2\sigma(e p \rightarrow e |c\bar{c}\rangle_1 X)}{dW dQ^2}$$

The three partonic diagrams shown in figure 5 should be taken into account




Figure 5: Boson-gluon-gluon fusion

and those obtained by reversing the orientation of the charm quark line. 6 diagrams in total.

Their contribution can be written as a convolution over the two-gluons density for unit area in the proton with respect to the fractions of longitudinal momentum carried $\rho_g^{(2)}(\xi_1, \xi_2, \mu)$:

$$\frac{d^2\sigma_{2g}^{(LO)}(e p \rightarrow e |c\bar{c}\rangle_1 X)}{dW dQ^2} = \int d\xi_1 \int d\xi_2 \rho_g^{(2)}(\xi_1, \xi_2, \mu) \hat{F}_{2g}[\xi_1, \xi_2, Q^2, S, W, \alpha_s(\mu)] \quad (2)$$

where $\sqrt{S}$ is the available center of mass energy for the electron-proton collision (at HERA $\sqrt{S} \simeq 300$ GeV) and $\alpha_s(\mu)$ is the strong running coupling constant.

In the low density limit $\rho_g^{(2)}(\xi_1, \xi_2, \mu)$ can be written as a product of single-gluon densities (the branches of a partonic cascade evolve independently):

$$\rho_g^{(2)}(\xi_1, \xi_2, \mu) = \frac{9}{8\pi R^2} \rho_g(\xi_1, \mu) \rho_g(\xi_2, \mu)$$

where R is the radius characteristic of the region in the proton populated by gluons.

It's natural to identify R with the proton radius $R \simeq 1fm$ (gluons are uniformly distributed inside the proton), but in the so-called "hot spot" effects R can be smaller (e.g R could be the radius of a constituent quark $R < 0.5fm$).

It should be recalled that eq. (2) is infrared divergent because of soft singularities that occur in the limit $\xi_1 \rightarrow 0$ or $\xi_2 \rightarrow 0$.

However gluon recombination favours gluons that are close in the rapidity space (they have greater probability to overlap and interact). We make therefore the following change of variables

$$\xi_1 \rightarrow \xi, \quad \xi_2 \rightarrow e^{-\eta} \xi$$

(η is the difference in rapidity for the gluon pair).

Then eq. (2) becomes

$$\frac{d^2 \sigma_{2g}^{(LO)}(e p \rightarrow e |c\bar{c}\rangle_1 X)}{dW dQ^2} = \frac{9}{8\pi R^2} \int_{-\infty}^{+\infty} d\eta e^{-\eta} \int_{\xi_{min}(\eta)}^{\xi_{max}(\eta)} d\xi \xi \rho_g(\xi, \mu)$$

$$\rho_g(e^{-\eta} \xi, \mu) \mathcal{C}(|\eta|) \hat{F}_{2g}[\xi, \eta, Q^2, S, W, \alpha_s(\mu)] \quad (3)$$

$\mathcal{C}(|\eta|)$ is here a correlation function which implements mathematically the previous guess, while

$$\xi_{min}(\eta) = \frac{Q^2 + 4m_c^2}{(1 + e^{-\eta}) Q^2} x, \quad \xi_{max}(\eta) = \frac{1}{1 + e^{-\eta}}$$

Let us assume now that

$$\mathcal{C}(|\eta|) = \delta(\eta)$$

This is, of course, a very strong hypothesis; it means that we allow recombination only for gluons with the same rapidity. Then one gets

$$\frac{d^2\sigma_{2g}^{(LO)}(e p \rightarrow e |c\bar{c}\rangle_1 X)}{dW dQ^2} = \frac{9}{8\pi R^2} \int_{\xi_{min}(0)}^{\xi_{max}(0)} d\xi \xi \rho_g^2(\xi, \mu) \hat{F}_{2g}[\xi, \eta = 0, Q^2, S, W, \alpha_s(\mu)] \quad (4)$$

Function $\hat{F}_{2g}[\xi, \eta = 0, Q^2, S, W, \alpha_s(\mu)]$ in eq. (4) has been computed in pQCD, neglecting the transverse components of incoming gluons momenta and their interactions (free gluon wave functions).

Its explicit expression is rather awkward, so we decided to put it in the appendix.

Eq. (4) has been numerically integrated by using gluon densities implemented in CERN **PDFLIB** library.

The result turns out to be infrared safe; this is due to the hypothesis made on $\mathcal{C}(|\eta|)$ shape which suppresses the initial state soft divergences.

Of course this contribution has to be summed to the corresponding one-gluon term $\frac{d^2\sigma_g^{(LO)}(e p \rightarrow e |c\bar{c}\rangle_1 X)}{dW dQ^2}$.

Therefore the leading order contribution to the hadronic differential cross section for $e p \rightarrow e J/\psi(z > z_0) X$ process can be written as follows

$$\begin{aligned} \frac{d^2\sigma^{(LO)}(e p \rightarrow e J/\psi(z > z_0) X)}{dW dQ^2} = & \frac{d^2\sigma^{(LO)}(e p \rightarrow e |c\bar{c}\rangle_1 X)}{dW dQ^2} \cdot \mathcal{H}(|c\bar{c}\rangle_1 \rightarrow J/\psi(z > z_0)) + \\ & \frac{d^2\sigma^{(LO)}(e p \rightarrow e |c\bar{c}\rangle_8 X)}{dW dQ^2} \cdot \mathcal{H}(|c\bar{c}\rangle_8 \rightarrow J/\psi(z > z_0)) \end{aligned} \quad (5)$$

with

$$\begin{aligned} \frac{d^2\sigma^{(LO)}(e p \rightarrow e |c\bar{c}\rangle_1 X)}{dW dQ^2} = & \frac{d^2\sigma_g^{(LO)}(e p \rightarrow e |c\bar{c}\rangle_1 X)}{dW dQ^2} + \frac{d^2\sigma_{2g}^{(LO)}(e p \rightarrow e |c\bar{c}\rangle_1 X)}{dW dQ^2} \end{aligned}$$

The term

$$\frac{d^2\sigma^{(LO)}(e p \rightarrow e |c\bar{c}\rangle_8 X)}{dW dQ^2}$$

is the Born term which contributes to the production of a colour octet $c\bar{c}$ pair and $\mathcal{H}\left(|c\bar{c}\rangle_{1,8} \rightarrow J/\psi(z > z_0)\right)$ stand for the fraction of partonic events of type

$$e p \rightarrow e |c\bar{c}\rangle_{1,8} X$$

which hadronize in a J/ψ vector meson with elasticity $z > z_0$.

As we have already mentioned $\frac{d^2\sigma_g^{(LO)}(e p \rightarrow e |c\bar{c}\rangle_1 X)}{dW dQ^2}$ is an infrared divergent quantity (soft singularities); on this subject, the next-to-leading order interference terms of the Born contribution with the virtual corrections for the $e g \rightarrow e c \bar{c}$ transition (which contribute to a colour octet $c\bar{c}$ pair production) should be considered.

In figure 6 are shown some of them as cut diagrams.

If we examine the colour structure of the interference terms we find that their contribution to the squared matrix element can be written as:

$$\overline{\sum} I(e g \rightarrow e c \bar{c}) = I_{QED} T(F) C_2(F) + I_{NA} C_2(A)$$

where I_{QED} , as before, takes contribution from “electrodynamical” diagrams (only quark-quark-gluon vertices), while I_{NA} contains the contributions of “non abelian” diagrams.

Notice that $\overline{\sum} I(e g \rightarrow e c \bar{c})$ has the same colour decomposition as $\overline{\sum} |M_{LO}(e g \rightarrow e g c \bar{c})|^2$; both contributions to $\frac{d^2\sigma_g^{(NLO)}(e p \rightarrow e c \bar{c} X)}{dW dQ^2}$ exhibit soft divergences; nevertheless the last quantity remains “infrared safe” due to a compensation mechanism.

This fact suggests to redefine $\frac{d^2\sigma_g^{(LO)}(e p \rightarrow e |c\bar{c}\rangle_1 X)}{dW dQ^2}$ by including part of the interference terms, which enter in $\frac{d^2\sigma_g^{(NLO)}(e p \rightarrow e |c\bar{c}\rangle_8 X)}{dW dQ^2}$, id est $\frac{d^2\tilde{\sigma}_{g,1}^{(LO)}(e p \rightarrow e c \bar{c} X)}{dW dQ^2}$ is defined by substituting

$$\overline{\sum} |M_{LO}(e g \rightarrow e |c\bar{c}\rangle_1 g)|^2 = S_{QED} C_1$$

with

$$\overline{\sum} \left| \tilde{M}_1^{(LO)}(e g \rightarrow e g c \bar{c}) \right|^2 \equiv (S_{QED} + I_{QED}) C_1$$

in the expression for $\frac{d^2\sigma_g^{(LO)}(e \rightarrow e c \bar{c})}{dW dQ^2}(X)$.
This operation has to be readsorbed in a redefinition of the hadroni-
zation fractions $\mathcal{H}\left(|c\bar{c}\rangle_{1,8} \rightarrow J/\psi(z > z_0)\right)$.




Figure 6: $e g \rightarrow e c \bar{c}$ interference terms

$\frac{d^2\sigma_{g,1}^{(LO)}(e \rightarrow e c \bar{c})}{dW dQ^2}(X)$ is found to be infrared safe in view of a Bloch-

Nordsieck compensation analogously to what happens in quantum electrodynamics; furthermore it can be obtained from the next-to-leading order contribution to $\frac{d^2\sigma_q(e \rightarrow e \ c \ \bar{c} \ X)}{dW dQ^2}$ (computed by Smith et al. [9] [10] [11]) simply by fixing $C_2(A) = 0$ and multiplying by the colour factor $\frac{1}{8}$.

Then we add the two-gluons contribution and we find that

$$\begin{aligned} & \frac{d^2\tilde{\sigma}_1^{(LO)}(e \ p \rightarrow e \ c \ \bar{c} \ X)}{dW dQ^2} = \\ & \frac{d^2\tilde{\sigma}_{g,1}^{(LO)}(e \ p \rightarrow e \ c \ \bar{c} \ X)}{dW dQ^2} + \frac{d^2\sigma_{2g}^{(LO)}(e \ p \rightarrow e \ |c\bar{c}\rangle_1 \ X)}{dW dQ^2} \end{aligned}$$

Now eq. (5) can be written as follows

$$\begin{aligned} & \frac{d^2\sigma^{(LO)}(e \ p \rightarrow e \ J/\psi(z > z_0) \ X)}{dW dQ^2} = \\ & \frac{d^2\tilde{\sigma}_1^{(LO)}(e \ p \rightarrow e \ c \ \bar{c} \ X)}{dW dQ^2} \cdot \tilde{\mathcal{H}}_1(c\bar{c} \rightarrow J/\psi(z > z_0)) + \\ & \frac{d^2\tilde{\sigma}_8^{(LO)}(e \ p \rightarrow e \ c \ \bar{c} \ X)}{dW dQ^2} \cdot \tilde{\mathcal{H}}_8(c\bar{c} \rightarrow J/\psi(z > z_0)) ; \quad (6) \end{aligned}$$

where

$$\frac{d^2\tilde{\sigma}_8^{(LO)}(e \ p \rightarrow e \ c \ \bar{c} \ X)}{dW dQ^2} = \frac{d^2\sigma^{(LO)}(e \ p \rightarrow e \ |c\bar{c}\rangle_8 \ X)}{dW dQ^2}$$

We have already pointed out that the soft divergences in the perturbative expansion for $\frac{d^2\sigma(e \rightarrow e \ |c\bar{c}\rangle_1 \ X)}{dW dQ^2}$ compensate, order by order,

with analogous singularities in $\frac{d^2\sigma(e \rightarrow e \ |c\bar{c}\rangle_8 \ X)}{dW dQ^2}$.

So the procedure introduced above, consisting in a subtraction of terms from $\frac{d^2\sigma(e \rightarrow e \ |c\bar{c}\rangle_8 \ X)}{dW dQ^2}$ to $\frac{d^2\sigma(e \rightarrow e \ |c\bar{c}\rangle_1 \ X)}{dW dQ^2}$, in order to compensate their soft divergences, can be extended, in principle, to all orders.

In this way are defined the higher order (“infrared safe”) corrections to $\frac{d^2\tilde{\sigma}_1(e \rightarrow e \ c \ \bar{c} \ X)}{dW dQ^2}$ and $\frac{d^2\tilde{\sigma}_8(e \rightarrow e \ c \ \bar{c} \ X)}{dW dQ^2}$.

Both these quantities depend, in general, on the renormalization scale μ (which we set equal to the factorization scale), while the

physical cross section $\frac{d^2\sigma(e\ p\rightarrow e\ J/\psi(z>z_0)\ X)}{dWdQ^2}$ does not depend on it.

As a consequence $\tilde{\mathcal{H}}_1$ and $\tilde{\mathcal{H}}_8$ have to depend in turns on μ in addition to z_0 , W , Q^2 and m_c .

In the following only z_0 dependence will be considered and all other effects will be neglected.

H1 collaboration has experimentally estimated the values of the differential cross sections $\frac{d\sigma(e\ p\rightarrow e\ J/\psi(z>0.2)\ X)}{dW}$ and $\frac{d\sigma(e\ p\rightarrow e\ J/\psi(z>0.2)\ X)}{dQ^2}$ for inelastic J/ψ electroproduction in $e^+ p$ collisions.

Therefore we have numerically integrated our leading order expression in eq. (6), with respect to Q^2 or W .

The factorization/renormalization scale is set as follows

$$\mu = \sqrt{Q^2 + 4m_c^2}$$

The right hand side of eq. (6) is then fitted to experimental data by using the minimization routines in **MINUIT** package of F.James and living $\tilde{\mathcal{H}}_1$, $\tilde{\mathcal{H}}_8$ and R as free parameters.

Different choices for the density of gluons are made, in order to compare the fit's results.

The following NLO gluon densities have been used: **MRS (G) (02.1995)**, **CTEQ 2pM**, **CTEQ 4M** and **GRV 94 HO**.



The results of these calculations are reported in table 1.

MRS (G) (02.1995) gluon density $R = 1.10 fm$, $\tilde{\mathcal{H}}_1(z_0=0.2) = 0.269$, $\tilde{\mathcal{H}}_8(z_0=0.2) = 2.75 \cdot 10^{-3}$ $\chi^2 = 6.74$, $d.o.f = 10 \Rightarrow \chi_{red}^2 = 0.674$
CTEQ 2pM gluon density $R = 0.442 fm$, $\tilde{\mathcal{H}}_1(z_0=0.2) = 0.175$, $\tilde{\mathcal{H}}_8(z_0=0.2) = 4.21 \cdot 10^{-3}$ $\chi^2 = 6.32$, $d.o.f = 10 \Rightarrow \chi_{red}^2 = 0.632$
CTEQ 4M gluon density $R = 0.393 fm$, $\tilde{\mathcal{H}}_1(z_0=0.2) = 0.109$, $\tilde{\mathcal{H}}_8(z_0=0.2) = 4.79 \cdot 10^{-3}$ $\chi^2 = 6.23$, $d.o.f = 10 \Rightarrow \chi_{red}^2 = 0.623$
GRV HO gluon density $R = 0.611 fm$, $\tilde{\mathcal{H}}_1(z_0=0.2) = 0.160$, $\tilde{\mathcal{H}}_8(z_0=0.2) = 5.32 \cdot 10^{-3}$ $\chi^2 = 6.51$, $d.o.f = 10 \Rightarrow \chi_{red}^2 = 0.651$



Table 1: Comparison of fit results for different gluon densities

In the following graphs H1 experimental points are compared with our results.



MRS (G) (02.1995) gluon density





CTEQ 2pM gluon density



CTEQ 4M gluon density



GRV HO gluon density



The errorbars are built-up summing in quadrature statistical and systematical errors (obviously χ^2 has been computed taking into account only statistical ones).

We remark that, even if these fits give values for $\tilde{\mathcal{H}}_8$ 1-2 orders of magnitude smaller than for $\tilde{\mathcal{H}}_1$, the perturbative terms connected to $\tilde{\mathcal{H}}_1$ are one order of magnitude smaller than those linked with $\tilde{\mathcal{H}}_8$; the latter contribution can be up to one order of magnitude smaller than that associated with $\tilde{\mathcal{H}}_1$.

The large discrepancies in the fitted parameters obtained using different gluon densities indicate that inelastic charmonium production provides a particularly sensitive tool to probe the gluonic structure of the hadrons; this is due to the fact that the two-gluons contribution, which depends on the square of the gluon density and then varies strongly with the choice of the density itself, is of the same order of magnitude as the single-gluon contribution (of the same perturbative order) in our kinematical range.

This fact suggests finally that important informations on the distribution of gluons in the transverse space in the proton could be obtained from hidden charm electroproduction.

Appendix

$$\begin{aligned}
\hat{F}_{2g} [\xi, \eta = 0, Q^2, S, W, \alpha_s(\mu)] &= \frac{\pi W}{2^4 3^2 \xi^5 S^2 Q^2 (Q^2 + W^2)^6} \{ \alpha^2 Q_c^2 \\
\alpha_s^2(\mu) \left\{ \frac{1}{m_c^2} \left\{ [Q^2 (2\xi - 1) + 2W^2 \xi] \sqrt{1 - \frac{4m_c^2}{Q^2 (2\xi - 1) + 2W^2 \xi}} \right. \right. \\
&\quad \left\{ 1920m_c^6 [2S^2 - 2S (Q^2 + W^2) + (Q^2 + W^2)^2] + \right. \\
&\quad 16m_c^4 \left\{ 4Q^2 [50S^2 - 50S (Q^2 + W^2) + 7 (Q^2 + W^2)^2] - \right. \\
&\quad 86 (Q^2 + W^2) [2S^2 - 2S (Q^2 + W^2) + (Q^2 + W^2)^2] \xi \} + \\
&\quad [2S^2 - 2S (Q^2 + W^2) + (Q^2 + W^2)^2] [Q^2 (2\xi - 1) + 2W^2 \xi] \\
&\quad [32Q^4 - 40Q^2 (Q^2 + W^2) \xi + 60 (Q^2 + W^2)^2 \xi^2] + 2m_c^2 \\
&\quad \left. \left. \left. \left. -32Q^4 [-16S^2 + 16S (Q^2 + W^2) + (Q^2 + W^2)^2] + 960SQ^2 \right. \right. \right. \\
&\quad \left. \left. \left. (Q^2 + W^2) (Q^2 + W^2 - S) \xi + 44 (Q^2 + W^2)^2 \xi^2 \right. \right. \right. \\
&\quad \left. \left. \left. [2S^2 - 2S (Q^2 + W^2) + (Q^2 + W^2)^2] \right\} \right\} \right\} + 12 \{ 320m_c^6 \\
&\quad [2S^2 - 2S (Q^2 + W^2) + (Q^2 + W^2)^2] + 8m_c^4 \{ 16Q^2 \\
&\quad [5S^2 - 5S (Q^2 + W^2) + (Q^2 + W^2)^2] - 42 (Q^2 + W^2) \\
&\quad [2S^2 - 2S (Q^2 + W^2) + (Q^2 + W^2)^2] \xi \} + 2SQ^2 (S - Q^2 - W^2) \\
&\quad [32Q^4 - 112Q^2 (Q^2 + W^2) \xi + 100 (Q^2 + W^2)^2 \xi^2] + m_c^2 \\
&\quad \{ 256SQ^4 (S - Q^2 - W^2) - 56Q^2 (Q^2 + W^2) [10S^2 - \\
&\quad 10S (Q^2 + W^2) + (Q^2 + W^2)^2] \xi + 84 (Q^2 + W^2)^2 [(Q^2 + W^2)^2 + \\
&\quad 2S^2 - 2S (Q^2 + W^2)] \xi^2 \} \left\{ \log \left\{ \frac{\left[1 + \sqrt{1 - \frac{4m_c^2}{Q^2 (2\xi - 1) + 2W^2 \xi}} \right]}{\left[1 - \sqrt{1 - \frac{4m_c^2}{Q^2 (2\xi - 1) + 2W^2 \xi}} \right]} \right\} \right\} \right\} \right\} \\
&\quad (7)
\end{aligned}$$

where $\alpha \simeq \frac{1}{137}$ is the e.m fine structure constant and $Q_c = \frac{2}{3}$ represents the quark charm electric charge (in e units).

References

- [1] H1 Collab., S.Aid et al., “*Elastic and Inelastic Photoproduction of J/ψ Mesons at HERA*” , Nucl.Phys. B472(1996) 3, 03/96
- [2] Zeus Collab., J.Breitweg et al., “*Measurement of Elastic J/ψ Photoproduction at HERA*” , Z.Phys. C75(1997) 215
- [3] Zeus Collab., J.Breitweg et al., “*Measurement of inelastic J/ψ Photoproduction at HERA*” , Z.Phys C76(1997) 599
- [4] H1 Collab., C.Adloff et al., “*Inelastic Photoproduction of J/ψ Mesons at HERA*” , Eur.Phys.J. C25 (2002) 1,25-39, 05/2002
- [5] H1 Collab., C.Adloff et al., “*Charmonium Production in Deep Inelastic Scattering at HERA*” , Eur.Phys.J. C10 (1999) 373-393, 03/99
- [6] S.J.Brodsky et al., “*Diffraction leptoproduction of vector mesons in QCD*” , Phys.Rev. D50(1994) 3134
- [7] L.Frankfurt, W.Koepf, M.Strikman, “*Hard diffractive electroproduction of vector mesons in QCD*” , Phys.Rev. D54(1996) 3194
- [8] L.Frankfurt, W.Koepf, M.Strikman, “*Diffraction heavy quarkonium photoproduction and electroproduction in QCD*” , Phys.Rev. D57(1998) 512
- [9] E.Laenen, S.Riemersma, J.Smith, W.L.van Neerven, “ *$O(\alpha_s)$ corrections to heavy flavour inclusive distributions in electroproduction*” , Nucl.Phys.B392: 229-250, 1993
- [10] E.Laenen, S.Riemersma, J.Smith, W.L.van Neerven, “*Complete $o(\alpha_s)$ corrections to heavy flavour structure functions in electroproduction*” , Nucl.Phys.B392: 162-228, 1993
- [11] B.W.Harris, J.Smith, “*Heavy quark correlations in deep inelastic electroproduction*” , Nucl.Phys.B452: 109-160, 1995